

Trefoil Factor 3 antibody

Cat. No. GTX134098

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application	WB, IHC-P
Reactivity	Human, Mouse

Package 100 μl, 25 μl

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 10 kDa. (Note)

PROPERTIES	
Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.19 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the C-terminus region of human Trefoil Factor 3. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 04 / 28 Page 1 of 2



DATA IMAGES



GTX134098 WB Image

Various whole cell extracts (30 μ g) were separated by 15% SDS-PAGE, and the membrane was blotted with Trefoil Factor 3 antibody (GTX134098) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX134098 IHC-P Image

Trefoil Factor 3 antibody detects secreted Trefoil Factor 3 protein by immunohistochemical analysis. Sample: Paraffin-embedded mouse intestine.

Trefoil Factor 3 stained by Trefoil Factor 3 antibody (GTX134098) diluted at 1:1500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX134098 IHC-P Image

Trefoil Factor 3 antibody detects Trefoil Factor 3 protein at cell membrane and cytoplasm by immunohistochemical analysis.

Sample: Paraffin-embedded mouse colon.

Trefoil Factor 3 stained by Trefoil Factor 3 antibody (GTX134098) diluted at 1:1500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 04 / 28 Page 2 of 2

€ 886-3-6208988 🔓 886-3-6208989 🐷 infoasia@genetex.com