

EAAT4 antibody

Cat. No. GTX134100

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	ICC/IF, IHC-P, IHC-Fr
Reactivity	Mouse, Rat

Package 100 μl, 25 μl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
IHC-Fr	1:100-1:1000

Not tested in other applications.

Properties	
Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the N-terminus region of mouse EAAT4. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 13 Page 1 of 2



DATA IMAGES



GTX134100 ICC/IF Image


EAAT4 antibody detects EAAT4 protein by immunofluorescent analysis.

Sample: DIV10 rat E18 primary cortical neuron cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: EAAT4 stained by EAAT4 antibody (GTX134100) diluted at 1:500.

Red: alpha Tubulin, stained by alpha Tubulin antibody (GT1141 (GTX628802) diluted at 1:500.

Red: alpha Tubulin, stained by alpha Tubulin antibody [GT114] (GTX628802) diluted at 1:500. Blue: Fluoroshield with DAPI (GTX30920).



GTX134100 IHC-Fr Image

EAAT4 antibody detects EAAT4 protein by immunohistochemical analysis. Sample: Frozen-sectioned mouse cerebellum.

Green: EAAT4 stained by EAAT4 antibody (GTX134100) diluted at 1:250.

Red: NF-H, stained by NF-H antibody [GT114] (GTX634289) diluted at 1:500.

Blue: Fluoroshield with DAPI (GTX30920).



GTX134100 IHC-P Image

EAAT4 antibody detects EAAT4 protein at cell membrane by immunohistochemical analysis. Sample: Paraffin-embedded rat brain.

EAAT4 stained by EAAT4 antibody (GTX134100) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 13 Page 2 of 2