

HuC antibody

Cat. No. GTX134128

| | |
|--------------|---------------------------|
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Applications | WB, ICC/IF, IHC-P, IHC-Fr |
| Reactivity | Mouse, Rat |

References (2)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | 1:500-1:3000 |
| ICC/IF | 1:100-1:1000 |
| IHC-P | 1:100-1:1000 |
| IHC-Fr | 1:100-1:1000 |

Not tested in other applications.

Calculated MW 40 kDa. ([Note](#))

Properties

| | |
|---------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Form | Liquid |
| Buffer | PBS, 20% Glycerol |
| Preservative | 0.025% ProClin 300 |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| Concentration | 1.26 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Carrier-protein conjugated synthetic peptide encompassing a sequence within the N-terminus region of mouse HuC. The exact sequence is proprietary. |
| Purification | Purified by antigen-affinity chromatography. |
| Conjugation | Unconjugated |




For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 10 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.


DATA IMAGES**GTX134128 IHC-Fr Image**

HuC antibody detects HuC protein by immunohistochemical analysis.

Sample: Frozen-sectioned mouse hippocampus.

Green: HuC stained by HuC antibody (GTX134128) diluted at 1:250.

Blue: Fluoroshield with DAPI (GTX30920).


**GTX134128 IHC-P Image**

HuC antibody detects HuC protein at cytoplasm and nucleus by immunohistochemical analysis.

Sample: Paraffin-embedded rat brain.

HuC stained by HuC antibody (GTX134128) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 10 min


**GTX134128 IHC-P Image**

HuC antibody detects HuC protein at cytoplasm and nucleus by immunohistochemical analysis.

Sample: Paraffin-embedded mouse brain.

HuC stained by HuC antibody (GTX134128) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 10 min

**GTX134128 IHC-P Image**

HuC antibody detects HuC protein at cytoplasm and nucleus by immunohistochemical analysis.

Sample: Paraffin-embedded mouse brain.

HuC stained by HuC antibody (GTX134128) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 10 min



For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 10 Page 2 of 2