


SARS-CoV-2 (COVID-19) Spike S1 Protein, B.1.1.7 / Alpha variant, His tag (active)

Cat. No. GTX136085-pro

| Applications | Binding Assay, WB | Package 100 μg |
|--------------|--------------------|-------------------|
| Species | SARS Coronavirus 2 | |

| Properties | | |
|-------------------|---|--|
| Form | Liquid | |
| Buffer | PBS | |
| Preservative | No preservatives | |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. For long-term storage after reconstitution, aliquot and store at -70°C or below. Do not vortex. | |
| Concentration | 1.1 mg/ml (Please refer to the vial label for the specific concentration.) | |
| Region/Sequence | SARS-CoV-2 Spike of QHD43416.1 (1-685 a.a) with S1-S2 cleavage site mutations (682-685GSAS) and variant mutatiosn (deletion 69-70, deletion 144, N501Y, A570D, D614G, P681H) and His tag at the C-terminus | |
| Expression System | HEK293 cells | |
| Purity | >95% | |
| Conjugation | Unconjugated | |
| Note | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. | |
| Note | Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product. | |

DATA IMAGES




GTX136085-pro Image

 $1~\mu g$ of GTX136085-pro SARS-CoV-2 (COVID-19) Spike S1 (del69-70, del44, N501Y, A570D, D614G...) Protein, His tag (active) protein was analyzed using SDS-PAGE and stained with coomassie blue and captured by monochrome camera.




For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 16 Page 1 of 2




GTX136085-pro Binding Assay Image

Functional ELISA analysis of immobilized recombinant Spike S1 protein(s) derived from different strains of SARS-CoV-2 virus (ie., Wild type; B.1.1.7 alpha variant) (coated at 2 μ g/mL) binding to soluble recombinant Human ACE2 (ECD) protein, mouse IgG Fc tag (active) (GTX135683-pro) (0.16-117.19 nM). Bound protein was detected by Goat Anti-Mouse IgG antibody (HRP) (GTX213111-01) (1:10000).




GTX136085-pro Binding Assay Image

Functional ELISA analysis of immobilized recombinant SARS-CoV-2 (COVID-19) Spike S1 Protein, B.1.1.7 / Alpha variant, His tag (active) (GTX136085-pro) (coated at 2 μ g/mL) binding to soluble recombinant Human ACE2 (ECD) protein, mouse IgG Fc tag (active) (GTX135683-pro) (0.16-117.19 nM). Bound protein was detected by Goat Anti-Mouse IgG antibody (HRP) (GTX213111-01) (1:10000).




GTX136085-pro Binding Assay Image

Inhibition analysis of immobilized recombinant SARS-CoV-2 (COVID-19) Spike S1 (del69-70, del44, N501Y, A570D, D614G...) Protein, His tag (active) (UK variant) (GTX136085-pro) and SARS-CoV-2 (COVID-19) Spike S1 protein, His tag (active) (GTX135817-pro) (coated at 2 μ g/mL) binding to soluble recombinant Human ACE2 (ECD) protein, mouse IgG Fc tag (active) (GTX135683-pro) (1000 ng/mL). ACE2 binding was inhibited by increasing concentrations of SARS-CoV-2 (COVID-19) Spike RBD antibody [HL1003] (GTX635792) (13.72-10000 ng/mL). Bound ACE2 was detected by Goat Anti-Mouse IgG antibody (HRP) (GTX213111-01) (1:10000).



GTX136085-pro WB Image

SARS-CoV-2 (COVID-19) Spike S1 (del69-70, del44, N501Y, A570D, D614G...) Protein, His tag (active) protein (GTX136085-pro, $0.5~\mu g$) was separated by 12% SDS-PAGE, and the membrane was blotted with the SARS-CoV-2 (COVID-19) Spike RBD antibody [HL257] (GTX635692) diluted at 1:5000.



GTX136085-pro Binding Assay Image

Inhibition analysis of immobilized recombinant SARS-CoV-2 (COVID-19) Spike S1 (del69-70, del44, N501Y, A570D, D614G...) Protein, His tag (active) (UK variant) (GTX136085-pro) and SARS-CoV-2 (COVID-19) Spike S1 protein, His tag (active) (GTX135817-pro) (coated at 2 μ g/mL) binding to soluble recombinant Human ACE2 (ECD) protein, mouse IgG Fc tag (active) (GTX135683-pro) (1000 ng/mL). ACE2 binding was inhibited by increasing concentrations of SARS-CoV-2 (COVID-19) Spike RBD antibody [HL1004] (GTX635793) (13.72-10000 ng/mL). Bound ACE2 was detected by Goat Anti-Mouse IgG antibody (HRP) (GTX213111-01) (1:10000).



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 16 Page 2 of 2