

# CD40 antibody [FGK4.5/ FGK45]

# Cat. No. GTX13944

Host	Rat
Clonality	Monoclonal
Isotype	lgG2a
Applications	WB, Activation
Reactivity	Mouse

Package 500 μg

## Applications

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
Activation	Assay dependent

Note: The FGK4.5 antibody is an agonistic antibody that has been shown to activate CD40 expressing APCs. FGK4.5 can also be used to inhibit CD40/CD154 interaction in vitro and in vivo.

Not tested in other applications.

Calculated MW 32 kDa. (Note)

Properties	
Form	Liquid
Buffer	Filter-sterilized PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant mouse CD40 fusion protein
Purification	Protein G purified From tissue culture supernatant
Purity	>95% (Determined by SDS-PAGE)
Endotoxin	< 0.002 EU/
Conjugation	Unconjugated



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 04 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

#### Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

## DATA IMAGES



#### GTX13944 WB Image

WB analysis of purified CD40 recombinant protein using GTX13944 CD40 antibody [FGK4.5/ FGK45].

Lane 1:1 ug reduced purified mouse CD40 with histidine & Fc tag at C-terminuso Lane 2:0.5 ug reduced purified mouse C040 with histidine & Fc tag at C-terminus

Dilution: 8 ug/ml



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 04 Page 2 of 2