

CD40 antibody [FGK4.5/ FGK45]

Cat. No. GTX13944

Host	Rat
Clonality	Monoclonal
Isotype	IgG2a
Applications	WB, Activation
Reactivity	Mouse

Package
500 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
Activation	Assay dependent

Note : The FGK4.5 antibody is an agonistic antibody that has been shown to activate CD40 expressing APCs. FGK4.5 can also be used to inhibit CD40/CD154 interaction in vitro and in vivo.

Not tested in other applications.

Calculated MW 32 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	Filter-sterilized PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant mouse CD40 fusion protein
Purification	Protein G purified From tissue culture supernatant
Purity	>95% (Determined by SDS-PAGE)
Endotoxin	< 0.002 EU/µg (Determined by LAL assay)
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES

**GTX13944 WB Image**

WB analysis of purified CD40 recombinant protein using GTX13944 CD40 antibody [FGK4.5/ FGK45].

Lane 1 : 1 ug reduced purified mouse CD40 with histidine & Fc tag at C-terminus

Lane 2 : 0.5 ug reduced purified mouse C040 with histidine & Fc tag at C-terminus

Dilution : 8 ug/ml



For full product information, images and publications, please visit our [website](#).