

CD19 antibody [1D3]

Cat. No. GTX14102

Host	Rat
Clonality	Monoclonal
Isotype	lgG2a
Application	FACS, Depletion, Neutralizing/Inhibition
Reactivity	Mouse

Package 500 μg

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
FACS	Assay dependent
Depletion	Assay dependent
Neutralizing/Inhibition	Assay dependent

Note: The 1D3 antibody has been reported to deplete B cells however, B cell depletion often requires treatment with a combination of 1D3, anti-mouse B220 (clone RA3.3A1/6.1), anti-mouse CD22 (clone Cy34.1), and anti-rat κ light chain (clone MAR 18.5) antibodies.

Not tested in other applications.

Calculated MW 60 kDa. (<u>Note</u>)

FormLiquidBufferFilter-sterilized PBSPreservativeNo preservativeStorageStore as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.ConcentrationBatch dependent (Please refer to the vial label for the specific concentration.)ImmunogenK562 cells expressing the extracellular domain of mouse CD19PurificationProtein G purified From tissue culture supernatantPurity>95% (Determined by SDS-PAGE)Endotoxin< 0.002 EU/*†g (Determined by LAL assay)	PROPERTIES	
PreservativeNo preservativeStorageStore as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.ConcentrationBatch dependent (Please refer to the vial label for the specific concentration.)ImmunogenK562 cells expressing the extracellular domain of mouse CD19PurificationProtein G purified From tissue culture supernatantPurity>95% (Determined by SDS-PAGE)Endotoxin< 0.002 EU/誊g (Determined by LAL assay)	Form	Liquid
Storage Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. Concentration Batch dependent (Please refer to the vial label for the specific concentration.) Immunogen K562 cells expressing the extracellular domain of mouse CD19 Purification Protein G purified From tissue culture supernatant Purity >95% (Determined by SDS-PAGE) Endotoxin < 0.002 EU/誊g (Determined by LAL assay)	Buffer	Filter-sterilized PBS
ConcentrationBatch dependent (Please refer to the vial label for the specific concentration.)ImmunogenK562 cells expressing the extracellular domain of mouse CD19PurificationProtein G purified From tissue culture supernatantPurity>95% (Determined by SDS-PAGE)Endotoxin< 0.002 EU/考g (Determined by LAL assay)	Preservative	No preservative
ImmunogenK562 cells expressing the extracellular domain of mouse CD19PurificationProtein G purified From tissue culture supernatantPurity>95% (Determined by SDS-PAGE)Endotoxin< 0.002 EU/眷g (Determined by LAL assay)	Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
PurificationProtein G purified From tissue culture supernatantPurity>95% (Determined by SDS-PAGE)Endotoxin< 0.002 EU/	Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Purity >95% (Determined by SDS-PAGE) Endotoxin < 0.002 EU/弮g (Determined by LAL assay)	Immunogen	K562 cells expressing the extracellular domain of mouse CD19
Endotoxin < 0.002 EU/眷g (Determined by LAL assay)	Purification	·
	Purity	>95% (Determined by SDS-PAGE)
Conjugation Unconjugated	Endotoxin	< 0.002 EU/弮g (Determined by LAL assay)
	Conjugation	Unconjugated



For full product information, images and publications, please visit our website.

Date 2024 / 05 / 05 Page 1 of 2



For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 05 Page 2 of 2