

GITR antibody [DTA-1]

Cat. No. GTX14350

| | |
|---------------------|------------|
| Host | Rat |
| Clonality | Monoclonal |
| Isotype | IgG2b |
| Applications | Activation |
| Reactivity | Mouse |

Package
500 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
|--------------------|----------------------|

| | |
|------------|-----------------|
| Activation | Assay dependent |
|------------|-----------------|

Note : The DTA-1 antibody is an agonistic antibody that is commonly used to induce GITR signaling in vivo.

Not tested in other applications.

Properties

| | |
|----------------------|--------------------------------------------------------------------------------------------------------|
| Form | Liquid |
| Buffer | Filter-sterilized PBS |
| Preservative | No preservatives |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. |
| Concentration | Batch dependent (Please refer to the vial label for the specific concentration.) |
| Immunogen | Mouse CD25+ CD4+ T cells |
| Purification | Protein G purified From tissue culture supernatant |
| Purity | >95% (Determined by SDS-PAGE) |
| Endotoxin | < 0.002 EU/µg (Determined by LAL assay) |
| Conjugation | Unconjugated |

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).