

## Transferrin antibody [F2H8G6]

Cat. No. GTX15790

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	ICC/IF, IP, ELISA, RIA
Reactivity	Human

Package  
100 µg

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	1:100-1:200
IP	Assay dependent
ELISA	1 µg/ml
RIA	Assay dependent

Not tested in other applications.

## Properties

Form	Liquid
Buffer	10mM Potassium Phosphate, 135mM NaCl, 2.5mM KCl, 0.1% BSA
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Human Transferrin
Purification	Protein A purified
Conjugation	Unconjugated


## Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

**DATA IMAGES**

**GTx15790 ELISA Image**

Sandwich ELISA analysis of recombinant transferrin protein ranging from 1.6-1000ng/ml using GTx15790 Transferrin antibody [F2H8G6] (detection antibody) at 1 µg/mL and transferrin chicken polyclonal antibody (capture antibody) at 1 µg/mL.


**GTx15790 ICC/IF Image**

ICC/IF analysis of HepG2 cells using GTx15790 Transferrin antibody [F2H8G6].

Fixation : Formalin

Permeabilization : 0.1% Triton X-100 in TBS for 10 minutes

Dilution : 1:100 overnight at 4 °C



For full product information, images and publications, please visit our [website](https://www.genetex.com).