

COX2 antibody, C-term

Cat. No. GTX15839

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, FCM
Reactivity	Human, Mouse

References (1)

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.1-0.3µg/ml
ICC/IF	10µg/ml
FCM	Assay dependent

Note : Strong expression of the protein seen in the cytoplasm and vesicles of HepG2 and NIH3T3 cells.

Not tested in other applications.

Calculated MW 69 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence C-NPTVLLKERSTEL, from the C Terminus of the protein sequence according to NP_000954.1.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX15839 FCM Image


FACS analysis of PFA fixed HeLa cells using GTX15839 COX2 antibody, C-term.

Blue : Primary antibody

Black : Isotype control

Permeabilization : 0.5% Triton

Dilution : 10µg/ml




GTX15839 WB Image

WB analysis of A549 (A) and Daudi (B) cell lysate using GTX15839 COX2 antibody, C-term.

Dilution : 0.1µg/ml

Loading : 35µg protein in RIPA buffer



GTX15839 ICC/IF Image

ICC/IF analysis of PFA fixed HepG2 cells using GTX15839 COX2 antibody, C-term.


Green : Primary antibody

Blue : DAPI

Negative control : Unimmunized goat IgG

Permeabilization : 0.15% Triton

Dilution : 10µg/ml



GTX15839 ICC/IF Image

ICC/IF analysis of PFA fixed NIH-3T3 cells using GTX15839 COX2 antibody, C-term.

Green : Primary antibody

Blue : DAPI

Negative control : Unimmunized goat IgG

Permeabilization : 0.15% Triton

Dilution : 10µg/ml



For full product information, images and publications, please visit our [website](#).