

Perforin antibody [CB5.4] (FITC)

Cat. No. GTX16075

| | |
|---------------------|-------------------------------|
| Host | Rat |
| Clonality | Monoclonal |
| Isotype | IgG2a |
| Applications | WB, ICC/IF, IHC-P, IHC-Fr, IP |
| Reactivity | Mouse |

References (3)

Package

50 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | 1 µg/mL |
| ICC/IF | Assay dependent |
| IHC-P | Assay dependent |
| IHC-Fr | Assay dependent |
| IP | Assay dependent |

Not tested in other applications.

Calculated MW 62 kDa. ([Note](#))

Properties

| | |
|----------------------|---|
| Form | Liquid |
| Buffer | PBS |
| Preservative | 0.02% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. Protect from light. |
| Concentration | Batch dependent (Please refer to the vial label for the specific concentration.) |
| Immunogen | Perforin (monoclonal CB5.4) antibody was raised against a recombinant mouse perforin, amino acids 98-534. |
| Purification | Purified IgG2a |
| Conjugation | Fluorescein isothiocyanate (FITC) Wavelength |



For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).