

DPAGT1 antibody

Cat. No. GTX16275

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, ELISA
Reactivity	Human, Mouse

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1 µg/mL
IHC-P	2.5 µg/mL
ELISA	Assay dependent

Not tested in other applications.

Calculated MW 46 kDa. ([Note](#))

Product Note At least four isoforms of DPAGT1 are known to exist; this antibody will recognize the two longest isoforms. DPAGT1 antibody is predicted to not cross-react with UHRF1BP1.

Properties

Form	Liquid
Buffer	PBS
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	DPAGT1 antibody was raised against a 17 amino acid synthetic peptide near the amino terminus of human DPAGT1. The immunogen is located within amino acids 30 - 80 of DPAGT1(NP_001373).
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

Date 2026 / 07 / 25 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES

GTX16275 IHC-P Image


IHC-P analysis of human kidney tissue using GTX16275 DPAGT1 antibody.

Working concentration : 20 µg/ml


GTX16275 IHC-P Image

IHC-P analysis of human kidney tissue using GTX16275 DPAGT1 antibody.

Working concentration : 2.5 µg/ml


GTX16275 WB Image

WB analysis of mouse kidney tissue lysate in (A) the absence and (B) the presence of blocking peptide using GTX16275 DPAGT1 antibody.

Working concentration : 1 µg/ml



For full product information, images and publications, please visit our [website](#).