

RAD23B antibody [5H1-A10-A7]

Cat. No. GTX16485

Host	Mouse
Clonality	Monoclonal
Isotype	lgG2b
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse, Rat, Hamster, Monkey

Package 100 μΙ

Applications

Application Note

Recommended Starting Dilutions:

For WB: Use at a dilution of 1:1000.

For ICC/IF: Use at a dilution of 1:100.

For IHC-P: Use at an assay depeendent dilution.

Not yet tested in other applications. Optimal dilutions should be determined experimentally by the researcher.

Calculated MW 43 kDa. (Note)

Properties	
Form	Liquid
Buffer	PBS, 0.5% BSA, 50% glycerol
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	Purified recombinant human RAD23B protein fragments expressed in E.coli.
Purification	Purified by affinity chromatography
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our website.

Date 2025 / 07 / 16 Page 1 of 2




DATA IMAGES




GTX16485 WB Image

WB analysis of various cell lysates using hHR23b antibody [5H1-A10-A7] at a dilution of 1:1000.




GTX16485 WB Image

WB analysis of various cell lysates using hHR23b antibody [5H1-A10-A7] at a dilution of 1:1000.



GTX16485 ICC/IF Image

ICC/IF analysis of HeLa cells using hHR23b antibody [5H1-A10-A7] at a dilution of 1:100.



GTX16485 IHC-P Image

IHC-P analysis of prostate cancer using hHR23b antibody [5H1-A10-A7] at a dilution of 1:100. Antigen retrieval was performed by pressure cooking in citrate buffer (pH 6.0).



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 07 / 16 Page 2 of 2

€ 886-3-6208988 📻 886-3-6208989 🐷 infoasia@genetex.com