

Cytokeratin 5 + 14 antibody [16.4]

Cat. No. GTX16570

| | |
|---------------------|----------------------------|
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype | IgG2a |
| Applications | WB, IHC-Fr, IHC (Formalin) |
| Reactivity | Human |

Package
100 µg

Applications

Application Note

IHC: Use at an assay dependent dilution. IHC-Fr: Use at an assay dependent dilution. WB: Use at an assay dependent dilution. Detects bands of approximately 58 and 50 kDa corresponding to cytokeratins 5 and 14 respectively. Optimal dilutions/concentrations should be determined by the end user.

Properties

| | |
|----------------------|--|
| Form | Liquid |
| Buffer | PBS |
| Preservative | 0.08% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| Concentration | 1 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Tissue / cell preparation (cytokeratin enriched extract of feline tongue epithelium). |
| Purification | IgG fraction |
| Conjugation | Unconjugated |

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).