

Aminoadipate aminotransferase antibody, N-term

Cat. No. GTX16585

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human

Package
100 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.2-2.5 µg/ml
IHC-P	2-10 µg/ml

Not tested in other applications.

Calculated MW 47 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	PBS, 2% Sucrose
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.5-1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	A synthetic peptide corresponding to a N-terminal region of Human Aminoadipate aminotransferase
Purification	Protein A purified
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 07 Page 1 of 2

DATA IMAGES

**GTX16585 IHC-P Image**

IHC-P analysis of human kidney tissue using GTX16585 Aminoadipate aminotransferase antibody at 4.0-8.0 μ g/ml.

70 kDa
60 kDa
48 kDa
36 kDa
21 kDa

GTX16585 WB Image

WB analysis of HepG2 cells using GTX16585 Aminoadipate aminotransferase antibody at 1.25 μ g/ml.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 07 Page 2 of 2