

# Orai1 antibody

## Cat. No. GTX16613

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, FCM, LCI
Reactivity	Mouse, Rat

References ( 4 )
Package
50 µl

## Applications

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
IHC-P	Assay dependent
FCM	Assay dependent
LCI	Assay dependent

Not tested in other applications.

Calculated MW 33 kDa. (Note)

**Product Note** We do not recommend use of this product for Human samples.

Properties	
Form	Liquid
Buffer	PBS, 1% BSA
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.65 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide (C)KFLPLKRQAGQPS, corresponding to amino acid residues 200-212 (2nd extracellular loop) of mouse Orai1 (Accession: Q8BWG9).
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 05 / 05 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

#### Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

#### DATA IMAGES




#### GTX16613 FCM Image

FACS analysis of WEHI cells using GTX16613 Orai1 antibody.

Black: Unstained cell

Green : Cell staining with primary antibody Antibody amount :  $10 \mu g$  antibody /  $0.5 \times 10^6$  cells




### GTX16613 LCI Image

Live cell imaging analysis of intact living RBL cells using GTX16613 Orai1 antibody.

Panel A: Primary antibody


Panel B: Merge of panel A with the live view of the cell

Dilution: 1:100



#### GTX16613 WB Image

WB analysis of rat spleen (lanes 1 and 4), rat pancreas (lanes 2 and 5), and mouse B-cell lymphoma (WEHI) (lanes 3 and 6) lysates using GTX16613 Orai1 antibody preincubated with or without immunogen peptide. Dilution: 1:200



## GTX16613 IHC-P Image

IHC-P analysis of rat pancreas tissue using GTX16613 Orai1 antibody. Orai1 (brown) is expressed in acinar cells. Hematoxilin is used as the counterstain.

Dilution: 1:50



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 05 / 05 Page 2 of 2