

Dab1 antibody

Cat. No. GTX16675

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IP
Reactivity	Mouse

Package
125 µg

Applications

Application Note

IP: Use at a concentration of 5 µg/ml. WB: Use at a concentration of 1 µg/ml. Detects a band of approximately 80 kDa (predicted molecular weight: 67 kDa). Optimal dilutions/concentrations should be determined by the end user.

Calculated MW 64 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS
Preservative	0.08% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recognizes Dab1 of 48 kDa, corresponding to amino acids 399-555 of mouse Dab1 GST fusion protein.
Purification	Protein G purified
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX16675 WB Image

GTX16675 at 1g/ml staining mouse DGTX1 (80kDa) in mouse brain extract by Western blot. Lane 1: GTX16674 (N terminal). Lane 2: GTX16675 (C terminal).



For full product information, images and publications, please visit our [website](#).