

# Src antibody [327]

**Cat. No. GTX16885**

<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1
<b>Applications</b>	WB, IP
<b>Reactivity</b>	Human, Mouse, Rat

**Package**  
50 µg

## Applications

### Application Note

For IP: Use at an assay dependent dilution. For WB: Use at a concentration of 2 -3 µg/ml. Predicted molecular weight: 59.7 kDa. GTX16885 may appear as a doublet on WB due to phosphorylation. This antibody can be used to precipitate active src kinase. Not suitable for IF, IHC-Fr or IHC-P. Optimal dilutions/concentrations should be determined by the researcher.

<b>Calculated MW</b>	61 kDa. ( <a href="#">Note</a> )
<b>Product Note</b>	Reacts with both v-Src and c-Src proteins.

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	50mM Sodium Phosphate, 0.2% Gelatin
<b>Preservative</b>	0.1% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
<b>Concentration</b>	0.1 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Antibody generated by immunizing BALB/c mice with full length native protein (purified) and then fusing with P3X63 Ag8.653 myeloma cells.
<b>Purification</b>	Purified IgG From hybridoma culture supernatant
<b>Conjugation</b>	Unconjugated

### Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).