

FBXL2 antibody, N-term

Cat. No. GTX17018

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Rat

References (1)

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.3-1µg/ml
IHC-P	5-10µg/ml

Note : Human Cerebral Cortex shows cytoplasm staining in neuronal bodies.

Not tested in other applications.

Calculated MW 47 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence VFSNNDEGLINKK-C, from the N Terminus of the protein sequence according to NP_036289.2.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 07 Page 1 of 2


DATA IMAGES

**GTX17018 WB Image**

WB analysis of human frontal cortex (A) and rat (B) brain lysate using GTX17018 FBXL2 antibody, N-term.

Dilution : 0.3 μ g/ml

Loading : 35 μ g protein in RIPA buffer

**GTX17018 IHC-P Image**

IHC-P analysis of human cerebral cortex using GTX17018 FBXL2 antibody, N-term.

Antigen retrieval : citrate buffer pH 6

Dilution : 10 μ g/ml



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 07 Page 2 of 2