

PDLIM1 antibody, C-term

Cat. No. GTX17022

Host	Goat	Package
Clonality	Polyclonal	100 µg
Isotype	IgG	
Applications	WB, IHC-P	
Reactivity	Human	

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.1-0.3µg/ml
IHC-P	5µg/ml

Not tested in other applications.

Calculated MW 36 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.50 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Peptide with sequence C-TPPEGYEVTVFPK, from the C Terminus of the protein sequence according to NP_066272.1.
Purification	Purified by ammonium sulphate precipitation followed by antigen affinity chromatography
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 13 Page 1 of 2

DATA IMAGES

**GTX17022 IHC-P Image**

IHC-P analysis of human placenta using GTX17022 PDLM1 antibody, C-term.

Antigen retrieval : citrate buffer pH 6

Dilution : 5µg/ml

250kDa
150kDa
100kDa
75kDa
50kDa
37kDa
25kDa
20kDa

GTX17022 WB Image

WB analysis of human lung lysate using GTX17022 PDLM1 antibody, C-term.

Dilution : 0.1µg/ml

Loading : 35µg protein in RIPA buffer



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 13 Page 2 of 2