

Apc6 antibody

Cat. No. GTX17065

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, ELISA
Reactivity	Human, Mouse

Package
100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1 - 2 µg/mL
IHC-P	5 µg/mL
ELISA	Assay dependent

Not tested in other applications.

Calculated MW 72 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	APC6 antibody was raised against an 18 amino acid synthetic peptide near the amino terminus of human APC6. The immunogen is located within the first 50 amino acids of APC6.
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated


Note For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).


DATA IMAGES

**GTX17065 IHC-P Image**

IHC-P analysis of human liver tissue using GTX17065 Apc6 antibody.
Working concentration : 5 µg/ml

**GTX17065 WB Image**

WB analysis of human liver tissue lysate using GTX17065 Apc6 antibody.
Working concentration : (A) 1 and (B) 2 µg/ml

**GTX17065 IHC-P Image**

IHC-P analysis of human liver tissue using GTX17065 Apc6 antibody.
Working concentration : 20 µg/ml



For full product information, images and publications, please visit our [website](#).