

Secretory Component Glycoprotein antibody [0.N.556]

Cat. No. GTX17377

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Application	ICC/IF, IHC-P
Reactivity	Human, Rat

Package
500 µl

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	Assay dependent
IHC-P	1:100-1:200 for 20 minutes at RT

Note : Antigen retrieval : Requires boiling tissue sections in 10mM citrate buffer, pH 6.0 for 10-20 minutes, followed by cooling at RT for 20 minutes

Not tested in other applications.

Product Note

Reacts with both free and polymeric Ig-bound secretory component. Shows no reaction in Western Blots of cell lines lacking secretory component.

PROPERTIES

Form	Liquid
Buffer	PBS, 0.2% BSA
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	Semi-purified human secretory component (SC).
Purification	Protein G purified
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES

GTX17377 IHC-P Image

IHC-P analysis of human colon carcinoma tissue using GTX17377 Secretory Component Glycoprotein antibody [0.N.556].



For full product information, images and publications, please visit our [website](#).