

## TOR1AIP1 antibody [RL13]

Cat. No. GTX17579

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	WB, ICC/IF, IHC-P, IP
Reactivity	Human, Mouse, Rat

Package  
200 µl

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
IHC-P	Assay dependent
IP	Assay dependent

Not tested in other applications.

**Calculated MW** 66 kDa. ( [Note](#) )

## Properties

Form	Liquid
Buffer	PBS
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	13.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Pore complex-lamina fraction isolated from rat liver nuclear envelopes.
Purification	Purified by PEG precipitation
Conjugation	Unconjugated


## Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

**DATA IMAGES**

**GTX17579 IHC-P Image**


IHC-P analysis of rat colon tissue using GTX17579 TOR1AIP1 antibody [RL13].

Right : Primary antibody

Left : a negative control without primary antibody

Dilution : 1:20 overnight

Antigen retrieval : 10mM sodium citrate followed by microwave treatment for 8-15 minutes.


**GTX17579 IHC-P Image**


IHC-P analysis of rat breast tissue using GTX17579 TOR1AIP1 antibody [RL13].

Right : Primary antibody


Left : a negative control without primary antibody

Dilution : 1:50 overnight

Antigen retrieval : 10mM sodium citrate followed by microwave treatment for 8-15 minutes.


**GTX17579 ICC/IF Image**

ICC/IF analysis of mouse NS-1 cells using GTX17579 TOR1AIP1 antibody [RL13].


**GTX17579 IHC-P Image**

IHC-P analysis of rat lymph node tissue using GTX17579 TOR1AIP1 antibody [RL13].

Right : Primary antibody

Left : a negative control without primary antibody

Dilution : 1:50 overnight

Antigen retrieval : 10mM sodium citrate followed by microwave treatment for 8-15 minutes.



For full product information, images and publications, please visit our [website](https://www.genetex.com).