

Human alpha Synuclein protein (monomer)

Cat. No. GTX17666-pro

Applications	Functional Assay	Package
Species	Human	200 µg, 100 µg

Applications

Application Note

Thioflavin T curve shows less β -sheet aggregation when Type 2 monomers (GTX17666-pro) are seeded with PFFs compared to Type 1 monomers (GTX17668-pro) seeded with PFFs.

Properties


Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Aliquot and store at -20°C or below. Avoid freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Region/Sequence	Full-length without tagged; MDVFMKGLSK AKEGWAAAE KTKQGVAAEA GKTKEGVLYV GSKTKEGVH GVATVAEKT EQVTNVGGAV VTGVTAVAQK TVEGAGSIAA ATGFVKKDQL GKNEEGAPQE GILEDMPVDP DNEAYEMPSE EGYQDYEPEA
Expression System	E. coli
Purification	Purified by ion-exchange chromatography
Purity	92 % by SDS-PAGE
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 19 Page 1 of 2


DATA IMAGES



GTX17666-pro Image

SDS-PAGE of ~14 kDa Human alpha Synuclein protein (active, monomer) (GTX17666-pro).

Lane 1: Molecular Weight Ladder (MW). Lane 2: BSA (2.5 µg). Lane 3: BSA (5 µg). Lane 4: Alpha Synuclein Protein Monomer (2.5 µg) (GTX17666-pro). Lane 5: Alpha Synuclein Protein Monomer (5 µg) (GTX17666-pro).



GTX17666-pro Image

Active alpha synuclein preformed fibrils (GTX17669-pro) seed the formation of new alpha synuclein fibrils from the pool of alpha synuclein monomers (GTX17668-pro). Thioflavin T is a fluorescent dye that binds to beta sheet-rich structures, such as those in alpha synuclein fibrils. Upon binding, the emission spectrum of the dye experiences a red-shift and increased fluorescence intensity. Thioflavin T emission curves show increased fluorescence (correlated to alpha synuclein protein aggregation) over time when 10 nM of active alpha synuclein preformed fibrils (GTX17669-pro) is combined with 100 µM of alpha synuclein monomer (GTX17668-pro), as compared to when 10 nM of control alpha synuclein preformed fibrils (GTX17667-pro) is combined with 100 µM of alpha synuclein monomer (GTX17668-pro) or 100 µM of alpha Synuclein monomer (GTX17666-pro). Thioflavin T ex = 450 nm, em = 485 nm.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 19 Page 2 of 2