

## Tropomyosin antibody [SPM224]

**Cat. No. GTX17784**

<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1
<b>Applications</b>	ICC/IF, IHC-P
<b>Reactivity</b>	Human, Rat, Chicken

References ( 1 )

Package

500 µl

## Applications

**Application Note**

For IF: Use at an assay dependent dilution. For IHC-P: Use at a dilution of 1:50. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10 min followed by cooling at RT for 20 min. Optimal dilutions/concentrations should be determined by the researcher.

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	PBS, 0.6% BSA
<b>Preservative</b>	0.09% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
<b>Concentration</b>	0.1 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Full length native protein (purified) (Chicken).
<b>Purification</b>	Protein G purified
<b>Conjugation</b>	Unconjugated

**Note**


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

## DATA IMAGES

**GTX17784 IHC-P Image**

Human heart stained with GTX17784 at 1/50 on paraffin sections



For full product information, images and publications, please visit our [website](#).