

Tropomyosin antibody [SPM224]

Cat. No. GTX17784

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	ICC/IF, IHC-P
Reactivity	Human, Rat, Chicken

References (1)

Package

500 µl

Applications

Application Note

For IF: Use at an assay dependent dilution. For IHC-P: Use at a dilution of 1:50. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10 min followed by cooling at RT for 20 min. Optimal dilutions/concentrations should be determined by the researcher.

Properties

Form	Liquid
Buffer	PBS, 0.6% BSA
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
Concentration	0.1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Full length native protein (purified) (Chicken).
Purification	Protein G purified
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX17784 IHC-P Image

Human heart stained with GTX17784 at 1/50 on paraffin sections



For full product information, images and publications, please visit our [website](#).