

FITC antibody

Cat. No. GTX19490

Host	Goat
Clonality	Polyclonal
Isotype	IgG F(ab') ₂
Applications	WB, ICC/IF, IHC-P, IHC-Fr, ELISA, IHC
Reactivity	Species independent

References (2)

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:200-1:1000
ICC/IF	Assay dependent
IHC-P	Assay dependent
IHC-Fr	Assay dependent
ELISA	1:2000-1:8000
IHC	Assay dependent

Not tested in other applications.

Product Note No reaction was observed against anti-Pepsin or anti-Goat IgG F(c).

Properties

Form	Liquid
Buffer	20mM Potassium Phosphate, 150mM NaCl
Preservative	0.01% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Fluorescein conjugated to Goat IgG
Purification	Purified by antigen-affinity chromatography. From serum
Conjugation	Unconjugated



For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).