

ICAM1 / CD54 antibody [15.2]

Cat. No. GTX20020

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	WB, ICC/IF, IHC-P, IHC-Fr, FCM, IP, ELISA, Blocking, Neutralizing/Inhibition
Reactivity	Human, Mouse, Rabbit, Pig

References (15)

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
IHC-P	Assay dependent
IHC-Fr	Assay dependent
FCM	≤ 1µg/10 ⁶ cells
IP	Assay dependent
ELISA	Assay dependent
Blocking	Assay dependent
Neutralizing/Inhibition	Assay dependent

Note : The suggested use of these reagents is in a final volume of 100 µL.

Not tested in other applications.

Calculated MW 58 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	Borate buffered saline
Preservative	No preservative
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
Concentration	0.1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Rheumatoid synovial cells and human monocytes



For full product information, images and publications, please visit our [website](#).

Purification	Purified IgG1
Conjugation	Unconjugated
Note	<p>For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.</p> <p>Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.</p>



For full product information, images and publications, please visit our [website](#).