

## ICAM1 / CD54 antibody [15.2]

Cat. No. GTX20020

<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1
<b>Applications</b>	WB, ICC/IF, IHC-P, IHC-Fr, FCM, IP, ELISA, Blocking, Neutralizing/Inhibition
<b>Reactivity</b>	Human, Mouse, Rabbit, Pig

References ( 15 )

Package

100 µg

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
IHC-P	Assay dependent
IHC-Fr	Assay dependent
FCM	≤ 1µg/10 <sup>6</sup> cells
IP	Assay dependent
ELISA	Assay dependent
Blocking	Assay dependent
Neutralizing/Inhibition	Assay dependent

**Note : The suggested use of these reagents is in a final volume of 100 µL.**

Not tested in other applications.

**Calculated MW** 58 kDa. ( [Note](#) )

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	Borate buffered saline
<b>Preservative</b>	No preservatives
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
<b>Concentration</b>	0.1 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Rheumatoid synovial cells and human monocytes



For full product information, images and publications, please visit our [website](#).

<b>Purification</b>	Purified IgG1
<b>Conjugation</b>	Unconjugated
<b>Note</b>	<p>For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.</p> <p>Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.</p>



For full product information, images and publications, please visit our [website](#).