

NCF1 antibody

Cat. No. GTX20795

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	WB
Reactivity	Human, Mouse, Rat, Rabbit, Bovine, Dolphin

Package

50 µg

Applications

Application Note

Western Blot - Approx 47kDa band seen in U937 cell lysates. Recommended for use at 0.2-2µg/ml.

Calculated MW 45 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	TBS, 0.5% BSA
Preservative	0.02% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Synthetic peptide: CSESTKRKLASAV.
Purification	IgG fraction Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTx20795 WB Image

GTx20795 staining (0.2g/ml) of U937 lysate (RIPA buffer, 30g total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



For full product information, images and publications, please visit our [website](#).