

Myc tag antibody (Agarose)

Cat. No. GTX21253

Host	Goat
Clonality	Polyclonal
Isotype	IgG
Applications	IP
Reactivity	Species independent

References (2) Package 100 μg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IP	Assay dependent

Note : Use 15-25 $\!\mu l$ of gel slurry per 0.1-1 mg of protein extract or lysate.

Not tested in other applications.

Properties	
Form	Liquid
Buffer	PBS
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	Amino acids 410-419 (EQKLISEEDL) of human myc conjugated to KLH.
Purification	Purified by antigen-affinity chromatography
Conjugation	Agarose
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 24 Page 1 of 1