Goat Anti-Rabbit IgG antibody (DyLight488)

Cat. No. GTX213110-04

Host	Goat
Clonality	Polyclonal
lsotype	lgG
Applications	WB, ICC/IF, IHC-P, IHC-Fr, IHC-Wm, FCM
Reactivity	Rabbit

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	1:100-1:2000
IHC-P	1:100-1:1000
IHC-Fr	1:100-1:1000
IHC-Wm	1:100-1:500
FCM	1:50-1:1000

Not tested in other applications.

Properties	
Form	Liquid
Buffer	PBS
Preservative	No preservative
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light.
Concentration	2 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Highly purified whole rabbit IgG
Purification	Purified by antigen-affinity chromatography.
Conjugation	DyLight488 Wavelength



For full product information, images and publications, please visit our <u>website</u>.

References (31) Package 500 µl



Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX213110-04 IHC-P Image

Double-labeled immunofluorescence photomicrographs of paraffin-embedded sections of mouse colon. Green: E-Cadherin antibody (GTX100443) diluted at 1:500. The signal was developed using goat anti-rabbit IgG antibody (Dylight488) (GTX213110-04).

Red: alpha Tubulin antibody [GT114] (GTX628802) diluted at 1:500. The signal was developed using goat anti-mouse IgG antibody (Dylight594) (GTX213111-05).

Blue: Fluoroshield with DAPI (GTX30920).

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min







Glypican 1 antibody [N3C3] (GTX104557) detects Glypican 1 protein by flow cytometry analysis. Sample: A431 cell.

Black: Unlabelled sample was used as a control.

Red: Glypican 1 antibody [N3C3] (GTX104557) dilution: 1:50.

Acquisition of 20,000 events were collected using the Rabbit IgG (DyLight488) (GTX113110-04) secondary antibody for FACS analysis.

GTX213110-04 IHC-Wm Image

Whole mount immunohistochemical analysis of paraformaldehyde-fixed 2 day-post-fertilization zebrafish embryo using Pax2a antibody (GTX128127) detected by anti-rabbit IgG antibody (Dylight488) (GTX213110-04). GTX128127 diluted at 1:100 and incubated overnight at 4°C. GTX213110-04 diluted at 1:500 and incubated 3 hours at room temperature.



GTX213110-04 IHC-Fr Image

Double-labeled immunofluorescence photomicrographs of frozen sections of mouse brain. Green: Vimentin antibody (GTX100619) diluted at 1:200. The signal was developed using goat anti-rabbit IgG antibody (Dylight488) (GTX213110-04). Red: ME1 antibody [GT15611] (GTX632190) diluted at 1:200. The signal was developed using goat anti-

mouse IgG antibody (Dylight594) (GTX213111-05).

Blue: Nuclear staining with Hoechst 33342.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 07 / 14 Page 2 of 2