# NFAT1 antibody [25A10.D6.D2]

# Cat. No. GTX22722

Host	Mouse
Clonality	Monoclonal
lsotype	lgG1
Application	WB, ICC/IF, IHC-P, IHC-Fr, FACS, IP, ChIP assay, Gel supershift assays
Reactivity	Human, Mouse, Rat

Reference (2) Package 100 μg

### APPLICATION

#### **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1 μg/ml
ICC/IF	1:100 - 1:1000
IHC-P	1:50-1:200
IHC-Fr	1:50-1:200
FACS	Assay dependent
IP	Assay dependent
ChIP assay	1-3 μl
Gel supershift assays	Assay dependent
Not tested in other applications.	

 Calculated MW
 100 kDa. (Note)

 Product Note
 This antibody does not cross react with NFAT2 (NFATc, NFATc1).

PROPERTIES	
Form	Liquid
Buffer	PBS
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Synthetic peptide corresponding to residues A(51) I S S P S G L A Y P D D V L D Y G L(69) of mouse NFATc2-A, B and C isoforms.




For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 18 Page 1 of 2




Purification	Protein G purified
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




#### GTX22722 IHC-P Image

IHC-P analysis of human spleen tissue using GTX22722 NFAT1 antibody [25A10.D6.D2]. Left : Primary antibody Right : Negative control without primary antibody Antigen retrieval : heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer, microwaved for 8-15 minutes Dilution : 1:100



## GTX22722 ICC/IF Image

ICC/IF analysis of HeLa cells using GTX22722 NFAT1 antibody [25A10.D6.D2]. Cells were probed without (right) or with(left) an antibody. Green : Primary antibody Blue : Nuclei Red : Actin Fixation : formaldehyde Dilution : 1:20 overnight at 4°C



#### GTX22722 ICC/IF Image

ICC/IF analysis of MCF-7 cells using GTX22722 NFAT1 antibody [25A10.D6.D2]. Cells were probed without (right) or with(left) an antibody. Green : Primary antibody Blue : Nuclei Red : Actin Fixation : formaldehyde Dilution : 1:20 overnight at 4°C



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 18 Page 2 of 2