

NFAT1 antibody [25A10.D6.D2]

Cat. No. GTX22722

| Host | Mouse |
|--------------|---|
| Clonality | Monoclonal |
| Isotype | lgG1 |
| Applications | WB, ICC/IF, IHC-P, IHC-Fr, FCM, IP, ChIP assay, Gel supershift assays |
| Reactivity | Human, Mouse, Rat |

References (2)
Package
100 μg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution | |
|-----------------------------------|----------------------|--|
| WB | 1 μg/ml | |
| ICC/IF | 1:100 - 1:1000 | |
| IHC-P | 1:50-1:200 | |
| IHC-Fr | 1:50-1:200 | |
| FCM | Assay dependent | |
| IP | Assay dependent | |
| ChIP assay | 1-3 μΙ | |
| Gel supershift assays | Assay dependent | |
| Not tested in other applications. | | |

| Calculated MW | 100 kDa. (<u>Note</u>) |
|---------------|---|
| Product Note | This antibody does not cross react with NFAT2 (NFATc1). |

| Properties | |
|---------------|--|
| Form | Liquid |
| Buffer | PBS |
| Preservative | 0.05% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| Concentration | 1 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Synthetic peptide corresponding to residues A(51) I S S P S G L A Y P D D V L D Y G L(69) of mouse NFATc2-A, B and C isoforms. |




For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 06 / 16 Page 1 of 2



| Purification | Protein G purified |
|--------------|---|
| Conjugation | Unconjugated |
| Note | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. |
| | Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product. |

DATA IMAGES



GTX22722 IHC-P Image

IHC-P analysis of human spleen tissue using GTX22722 NFAT1 antibody [25A10.D6.D2].


Left: Primary antibody

Right: Negative control without primary antibody

Antigen retrieval: heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer,

microwaved for 8-15 minutes

Dilution: 1:100



GTX22722 ICC/IF Image


ICC/IF analysis of HeLa cells using GTX22722 NFAT1 antibody [25A10.D6.D2]. Cells were probed without (right) or with(left) an antibody.

Green: Primary antibody

Blue: Nuclei Red: Actin

Fixation: formaldehyde

Dilution: 1:20 overnight at 4°C



GTX22722 ICC/IF Image

ICC/IF analysis of MCF-7 cells using GTX22722 NFAT1 antibody [25A10.D6.D2]. Cells were probed without

(right) or with(left) an antibody.

Green: Primary antibody

Blue: Nuclei Red: Actin

Fixation: formaldehyde

Dilution: 1:20 overnight at 4°C



For full product information, images and publications, please visit our website.

Date 2025 / 06 / 16 Page 2 of 2