

Cytohesin 2 antibody [10A12]

Cat. No. GTX22728

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	ICC/IF, IHC-P, IP
Reactivity	Human, Mouse

Package

100 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	1:20
IHC-P	1:25-1:200
IP	Assay dependent

Not tested in other applications.

Properties

Form	Liquid
Buffer	PBS, 0.1% BSA
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Purified, His-tagged cytohesin-2.
Purification	Purified by polyethylene glycol precipitation
Conjugation	Unconjugated

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 13 Page 1 of 2

DATA IMAGES

**GTx22728 IHC-P Image**


IHC-P analysis of human stomach tissue using GTx22728 Cytohesin 2 antibody [10A12].

Left : Primary antibody

Right : Negative control without primary antibody

Antigen retrieval : heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer, microwaved for 8-15 minutes

Dilution : 1:50

**GTx22728 IHC-P Image**


IHC-P analysis of human colon carcinoma tissue using GTx22728 Cytohesin 2 antibody [10A12].

Left : Primary antibody

Right : Negative control without primary antibody

Antigen retrieval : heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer, microwaved for 8-15 minutes

Dilution : 1:50

**GTx22728 ICC/IF Image**

ICC/IF analysis of HepG2 cells using GTx22728 Cytohesin 2 antibody [10A12]. Cells were probed without (right) or with(left) an antibody.


Green : Primary antibody

Blue : Nuclei

Red : Actin

Fixation : formaldehyde

Dilution : 1:100 overnight at 4 °C

**GTx22728 ICC/IF Image**

ICC/IF analysis of NIH-3T3 cells using GTx22728 Cytohesin 2 antibody [10A12]. Cells were probed without (right) or with(left) an antibody.

Green : Primary antibody

Blue : Nuclei

Red : Actin

Fixation : formaldehyde

Dilution : 1:100 overnight at 4 °C



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 13 Page 2 of 2