GRK1 antibody [G8]

Cat. No. GTX22775

Host	Mouse
Clonality	Monoclonal
lsotype	lgG1
Application	WB, ICC/IF, IHC-Fr, IP, IHC
Reactivity	Human, Mouse, Rat, Bovine, Dog, Chicken

Package 100 μg

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1 μg/ml
ICC/IF	2-3 μg/ml
IHC-Fr	Assay dependent
IP	Assay dependent
IHC	Assay dependent

Not tested in other applications.

Calculated MW	64 kDa. (<u>Note</u>)
Product Note	This antibody recognizes an epitope in the C-terminal region of GRK1a.

PROPERTIES	
Form	Liquid
Buffer	buffer
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Full length human GRK1.
Purification	Protein G purified
Conjugation	Unconjugated



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 19 Page 1 of 2




Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchas

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX22775 ICC/IF Image

ICC/IF analysis of HeLa cells using GTX22775 GRK1 antibody [G8]. Cells were probed without (left) or with(right) an antibody. Green : Primary antibody Blue : Nuclei Red : Actin Fixation : Formalin Permeabilization : 0.1% Triton X-100 in TBS for 5-10 minute Dilution : 1:200 and incubated overnight in a humidified chamber



IHC-Fr analysis of bovine retina tissue using GTX22775 GRK1 antibody [G8].



GTX22775 ICC/IF Image

ICC/IF analysis of murine cells using GTX22775 GRK1 antibody [G8]. Green : Primary antibody Blue : Nuclei Red : Actin Fixation : Formalin Permeabilization : 0.1% Triton X-100 in TBS for 5-10 minute Dilution : 1:200 and incubated overnight in a humidified chamber



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 19 Page 2 of 2