

## **VEGFA** antibody

## Cat. No. GTX22992

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	IHC-P
Reactivity	Human, Mouse, Rat

References ( 1 ) Package 500 µl

## Applications

## **Application Note**

Immunohistochemistry (formalin fixed paraffin embedded, rodent tissues only): use at a concentration of  $5 - 10\mu g$  / ml for 30 min at RT. Staining of formalin fixed tissues requires boiling tissue sections in 10mM Tris-HCl, pH 10.0, for 10 - 20 min followed by cooling at RT for 20 min. Neutralising (Int J Oncology, 8:505-511): assay dependent. Optimal dilutions / concentrations should be determined by the end user. Staining of formalin fixed tissues requires boiling tissue sections in 10mM Tris-HCl, pH 10.0, for 10 - 20 min followed by cooling at RT for 20 min.

Properties	
Form	Liquid
Buffer	PBS, 0.2% BSA
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
Concentration	0.2 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Human recombinant VEGF 165.
Purification	Protein A purified
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 07 Page 1 of 1