

Fibrillin 1 antibody [11C1.3]

Cat. No. GTx23090

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	WB, ICC/IF, IHC-P, IHC-Fr, EM
Reactivity	Human, Mouse, Bovine

References (6)

Package

500 μ l

Applications

Application Note

Electron Microscopy / Immunofluorescence / Immunohistochemistry: assay dependent. Western blot: use at a concentration of 1 - 2 μ g/ml for 2 hours at RT. By Western blot this detects a band of ~ 350 kDa, which corresponds to the predicted molecular weight of Fibrillin 1. Optimal dilutions/concentrations should be determined by the end user.

Calculated MW 312 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 0.2% BSA
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
Concentration	0.2 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Microfibrils from zonular apparatus of bovine eye.
Purification	Protein G purified
Conjugation	Unconjugated
For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 30 Page 1 of 1