

NOTCH1 antibody [A6]

Cat. No. GTX23294

Host	Mouse
Clonality	Monoclonal
Isotype	lgG2b
Applications	IHC-P, FCM
Reactivity	Human, Mouse

Package 500 μΙ

Applications

Application Note

For FACS: Use at an assay dependent dilution. For IHC-P: Use at 1/50 for 30 minute at room temperature. Staining of formalin fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10 - 20 min followed by cooling at room temperature for 20 min. Optimal dilutions / concentrations should be determined by the end user.

Product Note This antibody has not been tested for its cross reactivity with Notch 2.

Properties	
Form	Liquid
Buffer	PBS, 0.2% BSA
Preservative	0.1% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
Concentration	0.2 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encoding the ligand binding region of human Notch 1
Purification	Protein A affinity purified
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our website.

Date 2025 / 12 / 28 Page 1 of 1

€ 886-3-6208988 📻 886-3-6208989 🐷 infoasia@genetex.com