

Blood Group Lewis a antibody [T174]

Cat. No. GTx23356

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	IHC-P, IHC
Reactivity	Human

References (1)

Package

500 µl

Applications

Application Note

IHC-P: Use at a dilution of 1/40 or greater, for Biotin based detection systems. For optimal staining the primary antibody should be incubated 20-60 minutes at room temperature. Antigen retrieval is not essential. Optimal dilutions/concentrations should be determined by the end user.

Product Note

This antibody detects Lewis A (Type 1 chain) antigen and belongs to the BG-5 group of antibodies, one of a series of well-characterized murine monoclonal antibodies.

Properties

Form	Liquid
Buffer	PBS, 1% BSA
Preservative	0.1% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
Concentration	0.1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	SK-CO-10 colon cancer cell line (Human).
Purification	Protein G purified
Conjugation	Unconjugated
For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 07 Page 1 of 1