

CD99 antibody [O13]

Cat. No. GTX23382

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	IHC-P
Reactivity	Human

Package

500 µl

Applications

Application Note

IHC-P: Use at a dilution of 1/40 or greater for Biotin-based detection systems. For optimal staining, the primary antibody should be incubated 20-60 minutes at room temperature. Sections should be pretreated with an antigen unmasking solution. Optimal dilutions/concentrations should be determined by the end user.

Product Note This antibody is specific for the glycoprotein CD99, also referred to as p30/32^{MIC2} or the HBA71 antigen.

Properties

Form	Liquid
Buffer	PBS, 1% BSA
Preservative	0.1% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	MeWo human melanoma cells.
Purification	Protein A purified
Conjugation	Unconjugated

Note For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).