

Integrin beta 1 / CD29 (phospho Thr788/Thr789) antibody

Cat. No. GTX25189

Host	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Applications	WB, ICC/IF	
Reactivity	Human, Mouse, Cat	

References (1) Package 50 μl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:250
ICC/IF	1:250
ICC/IF	1:250

Not tested in other applications.

Calculated MW 88 kDa. (Note)

Properties	
Form	Liquid
Buffer	PBS, 0.1% BSA, 50% Glycerol
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	The antiserum was produced against a chemically synthesized phosphopeptide derived from the region of human integrin b1 receptor that contains threonines 788 and 789 (based on Swiss Protein database, accession number P05558). The sequence is conserved in human, mouse, rat and chicken.
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our website.

Date 2025 / 06 / 18 Page 1 of 2



DATA IMAGES



GTX25189 ICC/IF Image


ICC/IF analysis of THP-1 cells treated with 100ng/mL of IFN-gamma for 15 minutes using GTX25189 Integrin beta 1 / CD29 (phospho Thr788/Thr789) antibody. Panel e is untreated cell with no signal. Panel f represents control cells with no primary antibody to assess background.

Green: Primary antibody

Blue: Nuclei Red: Actin


Fixation: 4% paraformaldehyde

Permeabilization: 0.1% Triton X-100 for 10 minutes



GTX25189 WB Image

WB analysis of serum-starved mitotic HeLa cells generated by treatment with 100 ng/mL taxol for 16 hours using GTX25189 Integrin beta 1 / CD29 (phospho Thr788/Thr789) antibody. The data show that only the immunogen phosphopeptide blocks the signal, demonstrating the specificity of the antibody. The membrane treated with phosphatase (Lane 5) eliminates the signal further verifying that the antibody is phospho-specific.



GTX25189 WB Image

WB analysis of membrane enriched extracts (30 µg lysate) of HeLa (Lane 1) treated with Paclitaxel (Lane 2), HepG2 (Lane 3) treated with Paclitaxel (Lane 4), and 10 µL of conditioned media from HepG2 cell line (Lane 5) using GTX25189 Integrin beta 1 / CD29 (phospho Thr788/Thr789) antibody.

Dilution: 1:250



For full product information, images and publications, please visit our website.

Date 2025 / 06 / 18 Page 2 of 2