

c-Met (phospho Tyr1003) antibody

Cat. No. GTX25656

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, IHC
Reactivity	Human, Mouse, Dog, Hamster

Package 50 μl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
IHC-P	1:20-1:200
IHC	Assay dependent
Not tosted in other applications	

Not tested in other applications.

Calculated MW 156 kDa. (Note)

Properties	
Form	Liquid
Buffer	PBS, 0.1% BSA, 50% Glycerol
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	The antiserum was produced against a chemically synthesized phosphopeptide derived from the region of human c-Met that contains tyrosine 1003. The sequence is conserved in mouse and rat.
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 16 Page 1 of 2



DATA IMAGES



GTX25656 IHC-P Image


IHC-P analysis of mouse liver tissue using GTX25656 c-Met (phospho Tyr1003) antibody.

Right: Primary antibody

Left: Negative control without primary antibody

Antigen retrieval: 10mM sodium citrate (pH 6.0), microwaved for 8-15 min

Dilution: 1:100



GTX25656 WB Image

WB analysis of H69 cells unstimulated (Lane 1) or treated with 40 ng/mL HGF for 7.5 minutes (Lanes 2-5) using GTX25656 c-Met (phospho Tyr1003) antibody.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 16 Page 2 of 2