

Rabbit Anti-Goat IgG antibody, F(ab')<sub>2</sub> fragment, pre-adsorbed (TxRd)

Cat. No. GTX25751

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG F(ab') <sub>2</sub>
Application	ICC/IF, FACS, ELISA
Reactivity	Goat

Package  
250 µg

## APPLICATION

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	1:1000-1:5000
FACS	1:500-1:2500
ELISA	1:10000-1:50000

Not tested in other applications.

**Product Note** Pre-adsorbed with Human serum proteins. May react with immunoglobulins from other species.

## PROPERTIES

Form	Liquid
Buffer	10mM Sodium Phosphate, 250mM NaCl, 1% BSA
Preservative	0.01% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Goat IgG whole molecule
Purification	Purified by antigen-affinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorptions to remove any unwanted reactivities, pepsin digestion and chromatographic separation. From serum
Conjugation	Texas Red (TxRd) Ratio : 3.7 molecules TxRd per Rabbit IgG F(ab') <sub>2</sub> molecule.

For full product information, images and publications, please visit our [website](#).

**Note**

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).