

# PKC beta 2 (phospho Thr641) antibody

# Cat. No. GTX25785

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, FCM, IHC
Reactivity	Human, Rat

Package 50 μl

## Applications

#### **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	Assay dependent
FCM	1:20
IHC	Assay dependent
Not tosted in other applications	

Not tested in other applications.

Calculated MW 77 kDa. (Note)


Properties	
Form	Liquid
Buffer	PBS, 0.1% BSA, 50% Glycerol
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	The antiserum was produced against a chemically synthesized phosphopeptide derived from a region of human PKCbII that contains threonine 641. The sequence is conserved in mouse and rat.
Purification	Purified by antigen-affinity chromatography
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 15 Page 1 of 2

## DATA IMAGES



#### GTX25785 WB Image

WB (peptide competition) analysis of PMA-treated K652 cell lysates using GTX25785 PKC beta 2 (phospho Thr641) antibody with the non-phosphopeptide (Lane 2), a generic (pT) peptide (Lane 3), or the phosphopeptide immunogen (Lane 4). The data show that only the immunogen phosphopeptide blocks the signal, demonstrating the specificity of the antibody. The membrane treated with phosphatase (Lane 5) eliminates the signal further verifying that the antibody is phospho-specific.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 15 Page 2 of 2