

Goat Anti-Mouse IgG (F(ab')2) antibody, F(ab')2 fragment, pre-adsorbed (TxRd)

Cat. No. GTX25884

Host	Goat
Clonality	Polyclonal
Isotype	IgG F(ab')2
Applications	ICC/IF, FCM, Dot, ELISA
Reactivity	Mouse

Package

500 µg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	1:1000-1:5000
FCM	1:500-1:2500
Dot	Assay dependent
ELISA	1:10000-1:50000

Not tested in other applications.

Product Note	Pre-adsorbed with Bovine, Horse, Human, Rabbit, Rat and Sheep serum proteins. May react with immunoglobulins from other species.
--------------	--

Properties

Form	Liquid
Buffer	20mM Potassium Phosphate, 150mM NaCl, 1% BSA
Preservative	0.01% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Mouse IgG F(ab')2 fragment
Purification	Purified by antigen-affinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption to remove any unwanted reactivities, pepsin digestion and chromatographic separation. From serum
Conjugation	Texas Red (TxRd) Wavelength Ratio : 3.23 molecules TxRd per Goat IgG F(ab')2 molecule.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 09 Page 1 of 2

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 09 Page 2 of 2