

Goat Anti-Mouse IgM (mu chain) antibody, F(ab')<sub>2</sub> fragment (AP)**Cat. No. GTX25931**

<b>Host</b>	Goat
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG F(ab') <sub>2</sub>
<b>Applications</b>	WB, ELISA, IHC
<b>Reactivity</b>	Mouse

**Package**  
250 µg

## Applications

**Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:2000-1:20000
ELISA	1:10000-1:100000
IHC	1:500-1:2500

Not tested in other applications.

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	50mM Tris-HCl, 150mM NaCl, 1mM MgCl <sub>2</sub> , 0.1mM ZnCl <sub>2</sub> , 1% BSA, 50% Glycerol
<b>Preservative</b>	0.1% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE.
<b>Concentration</b>	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Mouse IgM whole molecule
<b>Purification</b>	IgG fraction This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgM coupled to agarose beads followed by solid phase adsorptions to remove any unwanted reactivities, pepsin digestion and chromatographic separation.
<b>Conjugation</b>	Alkaline Phosphatase (AP)
<b>Note</b>	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.  Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).