

Goat Anti-Rabbit IgG (Fc fragment) antibody, F(ab')2 fragment (AP)

Cat. No. GTX26024

| | | |
|--------------|----------------|---------|
| Host | Goat | Package |
| Clonality | Polyclonal | 250 µg |
| Isotype | IgG F(ab')2 | |
| Applications | WB, ELISA, IHC | |
| Reactivity | Rabbit | |

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | 1:500-1:2500 |
| ELISA | 1:2000-1:10000 |
| IHC | 1:200-1:1000 |

Not tested in other applications.

Properties

| | |
|---------------|--|
| Form | Liquid |
| Buffer | 50mM Tris-HCl, 150mM NaCl, 1mM MgCl ₂ , 0.1mM ZnCl ₂ , 1% BSA, 50% Glycerol |
| Preservative | 0.01% Sodium azide |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C. DO NOT FREEZE. |
| Concentration | 1 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Rabbit IgG Fc fragment |
| Purification | IgG fraction This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorptions to remove any unwanted reactivities, pepsin digestion and chromatographic separation. |
| Conjugation | Alkaline Phosphatase (AP) |
| Note | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. |



For full product information, images and publications, please visit our [website](#).

Date 2026 / 01 / 31 Page 1 of 1