

Goat Anti-Rabbit IgG (F(ab')<sub>2</sub>) antibody, F(ab')<sub>2</sub> fragment, pre-adsorbed (FITC)**Cat. No. GTX26108**

<b>Host</b>	Goat
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG F(ab') <sub>2</sub>
<b>Applications</b>	ICC/IF, FCM, Dot, ELISA
<b>Reactivity</b>	Rabbit

**Package**  
250 µg

## Applications

**Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	1:1000-1:5000
FCM	1:500-1:2500
Dot	Assay dependent
ELISA	1:10000-1:50000

Not tested in other applications.

**Product Note**

Pre-adsorbed with Bovine, Horse, Human, Mouse, Rat and Sheep serum proteins. May react with immunoglobulins from other species.

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	20mM Potassium Phosphate, 150mM NaCl, 1% BSA
<b>Preservative</b>	0.01% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. Protect from light.
<b>Concentration</b>	1 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Rabbit IgG F(ab') <sub>2</sub> fragment
<b>Purification</b>	Purified by antigen-affinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorptions to remove any unwanted reactivities, pepsin digestion and chromatographic separation. From serum
<b>Conjugation</b>	Fluorescein isothiocyanate (FITC) <a href="#">Wavelength</a>




For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

**Note**

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

## DATA IMAGES

**GTX26108 Dot Image**

Dot blot analysis of rabbit IgG using GTX26108 Goat Anti-Rabbit IgG (F(ab')<sub>2</sub>) antibody, F(ab')<sub>2</sub> fragment, pre-adsorbed (FITC).

Lane 1 : 50 ng

Lane 2 : 16.67 ng

Lane 3 : 5.56 ng

Lane 4 : 1.85 ng

Lane 5 : 0.62 ng

Dilution : 1:1000



For full product information, images and publications, please visit our [website](#).