

# CIKS antibody

**Cat. No. GTX26139**

|                     |            |
|---------------------|------------|
| <b>Host</b>         | Rabbit     |
| <b>Clonality</b>    | Polyclonal |
| <b>Isotype</b>      | IgG        |
| <b>Applications</b> | WB         |
| <b>Reactivity</b>   | Human      |

**Package**

50 µg

## Applications

### Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB                 | 0.5-1 µg/ml          |

Not tested in other applications.

**Calculated MW** 65 kDa. ( [Note](#) )

**Product Note** This antibody recognizes an epitope at the N-terminus of human CIKS/Act1 (Connection to IKK and SAPK/JNK).

## Properties

|                      |                                                                                                                                                                                                                            |
|----------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Form</b>          | Liquid                                                                                                                                                                                                                     |
| <b>Buffer</b>        | PBS                                                                                                                                                                                                                        |
| <b>Preservative</b>  | 0.02% Sodium azide                                                                                                                                                                                                         |
| <b>Storage</b>       | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| <b>Concentration</b> | Batch dependent (Please refer to the vial label for the specific concentration.)                                                                                                                                           |
| <b>Immunogen</b>     | Peptide corresponding to aa 2-15 of human CIKS.                                                                                                                                                                            |
| <b>Purification</b>  | Purified by affinity chromatography                                                                                                                                                                                        |
| <b>Conjugation</b>   | Unconjugated                                                                                                                                                                                                               |

### Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).